

Table 1. Estimated numbers of cases of HIV/AIDS, by year of diagnosis and selected characteristics, 2004–2007—34 states and 5 U.S. dependent areas with confidential name-based HIV infection reporting

	Year of diagnosis			
	2004	2005	2006	2007
Data for 34 states				
Age at diagnosis (yr)				
<13	212	189	169	159
13–14	41	40	45	40
15–19	1,081	1,216	1,409	1,703
20–24	3,714	3,875	4,184	4,907
25–29	4,524	4,547	4,884	5,771
30–34	5,353	5,024	4,686	5,089
35–39	6,359	5,907	5,678	6,088
40–44	6,011	5,889	6,003	6,554
45–49	4,286	4,338	4,377	5,172
50–54	2,645	2,698	2,862	3,489
55–59	1,473	1,531	1,512	1,938
60–64	771	729	741	942
≥65	696	657	643	803
Race/ethnicity				
American Indian/Alaska Native	177	180	163	228
Asian ^a	308	329	332	455
Black/African American	19,309	18,479	18,975	21,549
Hispanic/Latino ^b	6,183	6,383	6,590	7,484
Native Hawaiian/Other Pacific Islander	39	43	49	46
White	10,836	10,818	10,815	12,556
Transmission category				
Male adult or adolescent				
Male-to-male sexual contact	17,898	18,333	18,894	22,472
Injection drug use	3,198	2,990	2,931	3,133
Male-to-male sexual contact and injection drug use	1,413	1,308	1,195	1,260
High-risk heterosexual contact ^c	4,167	3,923	4,029	4,551
Other ^d	140	120	132	102
Subtotal	26,814	26,673	27,182	31,518
Female adult or adolescent				
Injection drug use	2,065	1,834	1,729	1,806
High-risk heterosexual contact ^c	7,967	7,852	8,033	9,076
Other ^d	103	90	80	96
Subtotal	10,135	9,775	9,842	10,977
Child (<13 yrs at diagnosis)				
Perinatal	177	162	134	139
Other ^e	37	30	36	20
Subtotal	214	192	170	159
Subtotal for 34 states	37,164	36,640	37,193	42,655
Data for U.S. dependent areas	1,234	1,391	1,338	1,429
Total^f	38,398	38,032	38,531	44,084

Note. These numbers do not represent reported case counts. Rather, these numbers are point estimates, which result from adjustments of reported case counts. The reported case counts have been adjusted for reporting delays and missing risk-factor information, but not for incomplete reporting. Data include persons with a diagnosis of HIV infection (not AIDS), a diagnosis of HIV infection and a later diagnosis of AIDS, or concurrent diagnoses of HIV infection and AIDS.

See Technical Notes for the list of areas that have had laws or regulations requiring confidential name-based HIV infection reporting since at least 2003.

^a Includes Asian/Pacific Islander legacy cases (see Technical Notes).

^b Hispanics/Latinos can be of any race.

^c Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^d Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.

^e Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

^f Includes persons of unknown race or multiple races and persons of unknown sex. Because column totals were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.